

Quistos da Rafe Mediana: Uma Raridade Inofensiva?

Median Raphe Cysts: A Harmless Rarity?

Ana Rita Curval^{1*}, Emília dos Santos², Elsa Marina Pereira Soares Loureiro²

***Autor Correspondente/Corresponding Author:**

Ana Rita Curval [rita_cur@hotmail.com]
Rua da Ponta 171, 4480-277 Junqueira,
Vila do Conde, Portugal
ORCID iD: 0000-0003-3084-0169

PALAVRAS-CHAVE: Criança; Doenças dos Genitais Masculinos/diagnóstico; Escroto; Períneo; Quistos/diagnóstico

KEYWORDS: Child; Cysts/diagnosis; Genital Diseases, Male/diagnosis; Perineum; Scrotum

A healthy 6-month-old boy was brought to a routine visit to the family physician (FF). The physical examination showed a thickened perineal raphe with several millimetric ovoid lesions from the scrotum to the perineum in the midline (Fig. 1). The remaining physical examination was unremarkable with no associated local inflammatory signs or other visible signs of anorectal malformations. Scrotal ultrasonography revealed retractable testicles, with no other findings. In his last visit to FF, at 18 months old, the lesions remained stable, with a slight increase in the thickening of the perineal raphe and small-sized cysts, located predominantly in the scrotal region (Fig. 2). The infant was referred to a Pediatric surgeon where the diagnosis of median raphe cysts (MRCs) was confirmed. Remained the indication for clinical surveillance.

MRCs are uncommon benign congenital lesions of the male genitalia, present anywhere from the perianal region to the scrotum and distal penis, being considered congenital defects in the embryologic

development.^{1,2} Diagnosis in children is rare, although probably underreported considering up to 75% are asymptomatic.¹ According to literature, most MRCs are found along the penile shaft and only a few cases report scrotum or perineal raphe cysts.³ Once they can mimic scrotal pearls in infants with anorectal malformations, a careful physical examination of the anogenital region should be performed.⁴ These cysts may grow with the child and clinical symptoms are more common in adulthood, due to infections^{1,4,5} or urinary symptoms like difficulty in voiding, hematuria, hematospermia, or sexual symptoms.^{1,6} Treatment for MRCs surgical excision in symptomatic patients. Conservative management could be considered for small asymptomatic lesions since histologic studies report no malignant potentiality.³ Spontaneous regression was barely reported.¹ It is important that pediatricians and FF recognize these lesions and their management to tranquilize and provide the appropriate information to parents to plan an adequate treatment.

1. Serviço de Pediatria, Centro Hospitalar Universitário de São João, Porto, Portugal. 2. USF Modivas, ACeS Grande Porto IV- Póvoa de Varzim/Vila do Conde, Vila do Conde, Portugal.

Recebido/Received: 2022/04/30 - Aceite/Accepted: 2023/05/10 - Publicado online/Published online: 2023/06/16 - Publicado/Published: 2023/09/30
© Author(s) (or their employer(s)) and Gazeta Médica 2023. Re-use permitted under CC BY-NC. No commercial re-use. © Autor (es) (ou seu (s) empregador (es)) e Gazeta Médica 2023. Reutilização permitida de acordo com CC BY-NC. Nenhuma reutilização comercial.



FIGURA 1. Millimetric ovoid lesions.

DECLARAÇÃO DE CONTRIBUIÇÃO/ CONTRIBUTORSHIP STATEMENT

ARC e ED: Revisão da literatura, escrita e revisão final do artigo

EL: Revisão do artigo

ARC and ED: Literature review, writing and final article review

EL: Article review

RESPONSABILIDADES ÉTICAS

CONFLITOS DE INTERESSE: Os autores declaram a inexistência de conflitos de interesse na realização do presente trabalho.

FONTES DE FINANCIAMENTO: Não existiram fontes externas de financiamento para a realização deste artigo.

CONFIDENCIALIDADE DOS DADOS: Os autores declaram ter seguido os protocolos da sua instituição acerca da publicação dos dados de doentes.

CONSENTIMENTO: Consentimento do doente para publicação obtido.

PROVENIÊNCIA E REVISÃO POR PARES: Não comissionado; revisão externa por pares.

ETHICAL DISCLOSURES

CONFLICTS OF INTEREST: The authors have no conflicts of interest to declare.

FINANCING SUPPORT: This work has not received any contribution, grant or scholarship.



FIGURA 2. Small-sized cysts.

CONFIDENTIALITY OF DATA: The authors declare that they have followed the protocols of their work center on the publication of data from patients.

PATIENT CONSENT: Consent for publication was obtained.

PROVENANCE AND PEER REVIEW: Not commissioned; externally peer reviewed.

REFERÊNCIAS

1. Krauel L, Tarrado X, Garcia-Aparicio L, Lerena J, Suñol M, Rodó J, et al. Median raphe cysts of the perineum in children. *Urology*. 2008;71:830-1. doi: 10.1016/j.urology.2007.11.131.
2. Patrizi A, Neri I, Lima M, Libri M, Gurioli C, Ravaioli GM. Congenital multiple median raphe cysts of the penis and scrotum. *J Paediatr Child Health*. 2019; 55:1389-90. doi: 10.1111/jpc.14534.
3. Shao IH, Chen TD, Shao HT, Chen HW. Male median raphe cysts: serial retrospective analysis and histopathological classification. *Diagn Pathol*. 2012;7:121. doi: 10.1186/1746-1596-7-121.
4. Soyer T, Karabulut AA, Boybeyi Ö, Günel YD. Scrotal pearl is not always a sign of anorectal malformation: median raphe cyst. *Turk J Pediatr*. 2013;55:665-6.
5. Lee JN, Kim HT, Cung SK: Median raphe cysts of the scrotum and perineum presenting with recurrent infection. *Korean J Urogenit Tract Infect Inflamm*. 2014;9:119-21.
6. Xambre L, Prisco R, Carreira F, Magalhães MF. Quistos do rafe médio - a propósito de um caso. *Acta Urol Port*. 2001;18:61-3.